

The annotated Catalogue of living Grylloblattida (insecta)

by

S. STOROZHENKO

After catalogue of GURNEY (1948) many articles were published on morphology, biology and systematics of the order *Grylloblattida*. Present catalogue summarizes new data on this interesting group of insects. Literature is cited according to YAMASAKI (1982a), abbreviations „a“, „b“ etc. agrees with place of articles on this bibliography (but CAUDELL, 1927 concern to CRAMPTON, 1927). Bibliography of YAMASAKI is complete and only recently published literature listed below.

In spite of the *Notoptera* CRAMPTON, 1915 is widely used for the order in recent time *Grylloblattida* E. WALKER (nom. trans.) must be preferred by the following aspects. After RASNITSYN (1976; 1980) it is clear that order consist of 3 fossil and recent taxa (*Paraplecoptera* + *Protoperlaria* + *Notoptera* or *Grylloblattodea*). The name *Notoptera* is very suitable because most of the fossil forms are well-winged, but some living orders of insects are apterous (for example *Mallophaga* etc.).

Order *Grylloblattida* general review (distribution, morphology, biology, bibliography):

COMSTOK, 1920: 268; IMMS, 1925: 318-320; CRAMPTON, 1927: 115-135; BRUES and MELANDER, 1932: 47-48; CRAMPTON, 1933: 127-166; WALKER, 1937: 1-10; GURNEY, 1937: 160-161; 1948: 90-98; 1953: 325-326; KAMP, 1953: 61; 1963: 53-68; BEY-BIENKO, 1966: 202-203; FUKUSHIMA, 1969a: 16-19, 1969b: 29-34; KAMP, 1970: 223-230; ASAHINA, 1971: 195-204; YAMASAKI, 1971: 1126; KAMP, 1979: 32-37; ANDO et al., 1979: 1-84; Mc E. KEVAN, 1979: 316-317; RENTZ, 1982: 1-18; NAMKUNG, 1982: 29-42; YAMASAKI, 1982a: 173-181; BROWN, 1982: 351-352; VICKERY and Mc E. KEVAN, 1983: 238-251.

Keys of genera and species: GURNEY, 1937: 160-161; 1953: 325-326; PRAVDIN and STOROZHENKO, 1977: 355-366; VICKERY and Mc E. KEVAN, 1983: 242.

Morphology

Head (with appendices): CRAMPTON, 1917a: 213-217; 1917c: 398-412; 1923: 77-107; 1926: 78-85; 1927: 123-125; WALKER, 1931: 519-522; 1933: 311-333; CRAMPTON, 1933: 133-140; KOYAMA, 1952: 54; FUKUSHIMA, 1967: 330; NAGASHIMA, 1982: 113-119.

Thorax (thoracic and cervical sclerites): CRAMPTON, 1915: 337-350; 1917: 402-404; 1926b: 214-216; 1927: 123-124; 1933: 140-145; WALKER, 1938:

589-600; FUKUSHIMA, 1967: 331; RÄHLE, 1970: 248-330; MATSUDA, 1970: 167-169.

Legs: CRAMPTON, 1927: 124-125; 1933: 145-147; WALKER, 1938: 591-600; FUKUSHIMA, 1967: 331-332.

Abdomen: WALKER, 1919a: 289-292; 1922: 39-41, 55; CRAMPTON, 1927: 125-127, 1933: 147-153; WALKER, 1943: 681-693; MATSUDA, 1976: 191-195; FUKUSHIMA, 1967: 331-332.

External genitalia and terminal segments of male: WALKER, 1922: 39-41, 55; EYER, 1924: 289-290; CRAMPTON, 1927: 126-127; 1933: 148-153; SNODGRASS, 1937: 19-22; WALKER, 1943: 688-693; 1956: 47-48; SNODGRASS, 1957: 20; SCUDDER, 1970: 55-56; MATSUDA, 1976: 192-194.

Ovipositor and terminal segments of female: CRAMPTON, 1917b: 225-237; WALKER, 1919a: 125-126; CRAMPTON, 1933: 126-150; WALKER, 1943: 684-688; MAHOTIN, 1952: 131-132; 1953: 37-41; WALKER, 1956: 48-49; SCUDDER, 1970: 57-58.

Sensilla: SLIFER, 1976: 275-279; PRITCHARD and SCHOLEFIELD, 1978: 208-210; EDWARDS and MANN, 1981: 177-188; EDWARDS and BALL, 1982: 220; McIVER and SUTCLIFFE, 1982: 137-157; BAKER, 1982: 341-344.

Eggs: FORD, 1926: 69-70; CRAMPTON, 1927: 134; 1933: 154; MATSUZAKI et al., 1979: 262-263; 1982: 85; ANDO and NAGASHIMA, 1982: 90-91.

Anatomy

Head: WALKER, 1931: 525-530; GOKAN et al., 1979: 258-271; 1982: 159-172; NAGASHIMA, 1982: 119-132.

Thorax: WALKER, 1938: 600-619; MATSUDA, 1970: 167-169.

Legs: WALKER, 1938: 600-619.

Abdomen: FORD, 1923: 207-319; WALKER, 1943: 693-701; MATSUDA, 1976: 191-195.

Organs of digestion: JUDD, 1948: 122-123; WALKER, 1949: 309-344; FUKUSHIMA, 1967: 333.

Heart and accessory structures: NUTTING, 1951: 501-597.

Neural system and its function: RAE and O'FARRELL, 1959: 76-83; FUKUSHIMA, 1967: 333; MORRISSEY and EDWARDS, 1979: 241-250; EDWARDS and MANN, 1981: 177-188.

Male reproductive organs: WALKER, 1943: 688-693; MATSUDA, 1976: 192-194; BACCETTI, 1981: 49-60; 1982: 71-78.

Female reproductive organs: CRAMPTON, 1933: 150-151; MATSUZAKI et al., 1979: 257-263; 1982: 79-85.

Spermatozoon: BACCETTI, 1982: 71-78.

Chromosome: NAKAMURA, 1946: 29-30.

Biology and Ecology

Habitat (general articles): WALKER, 1920: 139; FORD, 1926: 67-68; MILLS and PEPPER, 1937: 270-271; WALKER, 1937: 6-7; HISAUCHI, 1940: 48-50; GURNEY, 1948: 94-95; CHAPMAN, 1953: 39-41; KAMP, 1953: 62-63; FUKUSHIMA, 1965a: 76-79; 1965c: 96-97; 1966b: 36-38; STOROZHENKO, 1979: 18-21; RENTZ, 1982: 12-16; YAMASAKI, 1982c: 13-17.

Influence of temperature: BUCKEL, 1925: 35-36; FORD, 1926: 68; MILLS and PEPPER, 1937: 271-274; WALKER, 1937: 6; EDWARDS and NUTTING, 1950: 33-44; HENSON, 1957a: 637; FUKUSHIMA, 1965b: 34; EDWARDS, 1982: 24-25.

Behaviour: FORD, 1926: 67-68; YANO, 1927: 111; MILLS and PEPPER, 1937: 270-274; CAMPBELL, 1949: 1-5; FUKUSHIMA, 1966a: 33; PRITCHARD and SCHOLEFIELD, 1978: 205-208; NAGASHIMA et al., 1982: 46-47.

Nutrition: FORD, 1926: 68; ASAHINA, 1929: 195; PRITCHARD and SCHOLEFIELD, 1978: 205-208; EDWARDS, 1982: 22-23.

Life history: KENNEDY, 1928: 368; HENSON, 1957b: 149-152; NAGASHIMA et al., 1982: 43-59; VISSCHER et al., 1982: 68.

Copulation: FORD, 1926: 69; NAGASHIMA et al., 1982: 47-49; VISSCHER et al., 1982: 64-67.

Oviposition and number of eggs: FORD, 1926: 69-70; FUKUSHIMA, 1968: 28; NAGASHIMA et al., 1982: 49-51; VISSCHER et al., 1982: 67.

Embryogenesis: FUKUSHIMA, 1969d: 24; ANDO and NAGASHIMA, 1982: 89-95.

Environment and descriptions of larvae: SILVESTRI, 1927: 107-119; WALKER, 1937: 8; FUKUSHIMA, 1966b: 36-38; NAGASHIMA et al., 1982: 52-54; YAMASAKI, 1982b: 97-111.

Rearing techniques: FORD, 1926: 68; 1937: 3-27; FUKUSHIMA, 1965a: 76-79; 1966b: 36-38; NAGASHIMA et al., 1982: 43-44; VISSCHER et al., 1982: 62-63.

Color-form: EDWARDS, 1958: 49; FUKUSHIMA, 1969c: 7.

Affinities

The affinities of *Grylloblattida* are widely discussed in literature, but according to RASNITSYN (1976: 1980) it is clear what *Grylloblattidae* though *Blattogryllidae* connect with carboniferous-yurassic *Protoblattodea* + *Paraplecoptera* complex (about 28 families).

WALKER, 1914: 96-99; CRAMPTON, 1917b: 225-237; 1919: 47-48; WALKER, 1919a: 271-272; 1922: 58-64; CRAMPTON, 1926a: 83-84; 1926b: 240; 1927: 135; IMMS, 1927a: 57; 1927b: 36-39; SILVESTRI, 1927: 118-119; 1931: 293-294; CRAMPTON, 1933: 156-164; 1935: 97-111; WALKER, 1937: 2-5; 1949: 331-337; SNODGRASS, 1957: 20; BLACKITH and BLACKITH, 1968: 126; KRISTENSEN, 1973: 17-18; KAMP, 1973: 1235-1249; RASNITSYN, 1976: 502; HEMING, 1977: 80; McE-KEVAN, 1977a: 18, 20; RASNITSYN, 1980: 150-154; RENTZ, 1982: 2-3; VICKERY and McE-KEVAN, 1983: 238.

Taxonomy

Order *Grylloblattida* E. WALKER, 1914 (nom. trans. BEY-BIENKO, 1962) (= fossil *Paraplecoptera* + *Protoperlaria* + recent *Notoptera*)

Grylloblattida (pro order) BEY-BIENKO, 1962: 19; 1966: 202; ROHDENDORF, 1977: 20; PRAVDIN and STOROZHENKO, 1977: 352; RASNITSYN, 1980: 150; STOROZHENKO and OLIGER, 1984: 729.

Notoptera (pro order) CRAMPTON, 1915: 347; COMSTOK, 1920: 268; CAUDELL and KING, 1924: 60; CAUDELL, 1924: 92; KISHIDA, 1929: 104; McE-KEVAN, 1977b: 13; 1979: 316; VICKERY and McE-KEVAN, 1983: 238.

Notoptera (pro suborder *Orthoptera* s. str.) CRAMPTON, 1933: 160; 1935: 110.

Notopterina (pro order) DURDEN, 1969: 1159.

Grylloblattodea (pro order) IMMS, 1925: 318; BRUES and MELANDER, 1932: 47; KAMP, 1953: 61; FUKUSHIMA, 1969a: 16; MATSUDA, 1970: 167; ASAHINA, 1971: 195; KAMP, 1973: 1235; MATSUDA, 1976: 191; NAMKUNG, 1982: 29.

Grylloblattodea (pro suborder *Orthoptera* s. str.) NUTTING, 1951: 512.

Grylloblattodea (pro suborder *Notoptera*) McE-KEVAN, 1977b: 13; VICKERY and McE-KEVAN, 1983: 239.

Grylloblattoidea (pro order) BRUES and MELANDER, 1915: 1, 10, 13; WALKER, 1922: 39.

Grylloblattoidea (pro suborder *Orthoptera*) WALKER, 1919: 289; SILVESTRI, 1927: 119; SNODGRASS, 1937: 19.

Grylloblattidea (pro infraorder *Notoptera*) McE-KEVAN, 1977b: 13; VICKERY and McE-KEVAN, 1983: 239.

Grylloblattoidea (pro order) CRAMPTON, 1917c: 409.

Grylloblattoidea (pro order) CRAMPTON, 1920: 118.

Grylloblattaria (pro order) BRUNER, 1915, 195; JUDD, 1948: 96, 122; WALKER, 1949: 309; 1956: 47; BROWN, 1982: 351.

Grylloblattinae (pro suborder *Orthoptera* s. str.) BEIER, 1955: 31-304; BEIER, 1972: 2-217.

Archiorthoptera (pro order) CRAMPTON, 1933: 160.

Superfamily *Grylloblattoidea* E. WALKER, 1914 (nom. trans. RASNITSYN, 1976) (= fossil *Megakhosaridea*)

RASNITSYN, 1976: 503; McE-KEVAN, 1977b: 13; VICKERY and McE-KEVAN, 1983: 239.

Family *Grylloblattidae* E. WALKER, 1914.

WALKER, 1914: 93; COMSTOK, 1920: 268; CAUDELL and KING, 1924: 60; HANDLIRSCH, 1925: 445, 457-458; HEBARD, 1930: 381; SILVESTRI, 1931: 294; BRUES and MELANDER, 1932: 48; LAMEERE, 1935: 287-290; GURNEY, 1937: 159; CHOPARD, 1938: 35-380; GURNEY, 1948: 86; CHOPARD, 1949: 232-235; GURNEY, 1953: 325; BRUES et al., 1954: 94; GURNEY, 1961: 67; FUKUSHIMA, 1969a: 16; ASAHINA, 1971: 195; RASNITSYN, 1976: 502; McE-KEVAN, 1977b: 13; VICKERY and McE-KEVAN, 1983: 239.

Subfamily *Grylloblattinae* E. WALKER, 1914 (nom. trans. VICKERY et McE-KEVAN, 1983).

Grylloblattinae VICKERY and McE-KEVAN, 1983: 240, 241.

Galloisianinae VICKERY et McE-KEVAN, 1983: 240, syn. n. (The form of pronotum of some *Galloisiana* and *Grylloblattina* are similar with *Grylloblatta*, but the degrees of reduction of eyes and pulvilli of tarsi are not sufficient for erecting a subfamily).

1. Genus *Grylloblatta* E. WALKER, 1914.

WALKER, 1914: 93. Type species: *Grylloblatta campodeiformis* E. WALKER, by original designation.

SILVESTRI, 1927: 112 (part.); GURNEY, 1937: 160, 164; 1948: 86-95, 97; KAMP, 1963: 53; FUKUSHIMA, 1969a: 16; ASAHINA, 1971: 196, 201; KAMP, 1979: 27; RENTZ, 1982: 1; VICKERY and McE-KEVAN, 1983: 241.

11 species in North America (U.S.A. and Canada).

1. *Grylloblatta barberi* CAUDELL, 1924

CAUDELL, 1924b: 369 (Holotype: ♂ (larvae), U.S.A., California, Plumas County, North Fork of the Feather River at the junction of Butt Creek, in U.S. Nat. Mus., N 27265, Washington).

WALKER, 1937: 9; GURNEY, 1937: 160; 1948: 90, 98, fig. 11; 1953: 326, 330; KAMP, 1953: 61; 1963: 55, 65, figs. 1-7 (Plesiotype, ♀, in U.S. Nat. Mus., Washington); 1970: 225, fig. 1; RENTZ, 1982: 5, 7, 10.

Grylloblatta campodeiformis: CAUDELL, 1923a: 148-150; 1923b: 261.
Distribution. U.S.A.: California (Plumas County).

2. *Grylloblatta bifratrilecta* GURNEY, 1953.

GURNEY, 1953: 325, 328-330, figs. 4, 7, 15 (Holotype: ♂, U.S.A., California, Sonora Paß (about 8600-10000 foot), in U.S. Nat. Mus., Washington, N 61132, allotype: ♀, 2 miles W. Sonora Paß, in U.S. Nat. Mus.) KAMP, 1963: 65, fig. 43; 1970: 225, fig. 1; RENTZ, 1982: 5, 7, 10.
Distribution. U.S.A.: California (Tuolumne County).

3. *Grylloblatta campodeiformis* E. WALKER, 1914.

WALKER, 1914: 94, pl. IX, figs. 1-7. (Holotype: ♀, Canada, Sulphur Mountain, Banff (6500 foot) in R.O.M., Toronto).
WALKER, 1919b: 131, pl. XIII, figs. 1-7, pl. IX, figs. 8-15 (♂, larvae); COMSTOK, 1920: 268; CAUDELL, 1924b: 370; CAUDELL and KING, 1924: 58, figs. 4, 5; EYER, 1924: 289 (Australia, mistake); BUCKELL, 1925: 35-36; FORD, 1926: 66; IMMS, 1927: 57; SILVESTRI, 1927: 119, figs. XII (1-3); HEBARD, 1930: 381; SILVESTRI, 1931: 292-293, figs. 1-3; STRANDT, 1937: 38-39, fig. 7; MILLS and PEPPER, 1937: 269; WALKER, 1937: 8; GREGSON, 1938: 63-64; 1939: 29-30; SPENSER, 1945: 20 (part.); PLETSH, 1947: 17-20; GURNEY, 1948: 90, 97-98, figs. 3, 8-11, text-figs. A, C, D; SPENCER, 1952: 36; KAMP, 1953: 61; 1963: 65, fig. 43; VICKERY et al., 1974: 5; SLIFER, 1976: 275; PRITCHARD and SCHOLEFIELD, 1978: 205.

Grylloblatta campestris (lapsus calami): BEAMER, 1933: 236.
Divided into 3 subspecies.

3a. *Grylloblatta campodeiformis campodeiformis* E. WALKER, 1914.

GURNEY, 1937: 160; 1953: 325, 331, figs. 2, 3, 8, 9, 13; HENSON, 1957b: 149-152; CORNER, 1960: 49; GURNEY, 1961: 72; KAMP, 1970: 225, fig. 1; 1979: 27-28, figs. 19-28; RENTZ, 1982: 5, 6; VICKERY and McE-KEVAN, 1983: 242, figs. 80a, b, 82. (See also under *Grylloblatta campodeiformis*).
Distribution. Canada: S.E. British Columbia, S.W. Alberta; U.S.A.: N. Idaho, W. Washington.

3b. *Grylloblatta campodeiformis athapaska* KAMP, 1979.

KAMP, 1979: 27, figs. 1-9. (Holotype: ♂, Canada, British Columbia, Stone Provincial Park, Summit Lake, Mt. St. Paul, in Canadian National Collection, Ottawa; allotype: ♀, some locality and disposition).
RENTZ, 1982: 5, 12; VICKERY and McE-KEVAN, 1983: 242, 247.
Distribution. Canada: N.W. British Columbia.

3c. *Grylloblatta campodeiformis nahanni* KAMP, 1979.

KAMP, 1979: 28, figs. 10-18 (Holotype: ♂, Canada, British Columbia, Cassiar, Cassiar Mountain Range, Mt. McDane (1647 m), in Canadian National Collection, Ottawa; allotype: ♀, some locality (Limestone Peak, 1830 m), in University of British Columbia).
RENTZ, 1982: 5, 12; VICKERY and McE-KEVAN, 1983: 242, 248.
Distribution. Canada: N.E. British Columbia.

4. *Grylloblatta chandleri* KAMP, 1963.

KAMP, 1963: 57, figs. 8-22. (Holotype: ♂, U.S.A., California, Lassen County, Eagle Lake (5180 foot), in American Mus. Nat. Hist., New York; allotype: ♀, some locality, in U.S. Nat. Mus., Washington).
KAMP, 1970: 225, fig. 1; RENTZ, 1982: 5, 7, 10.
Grylloblatta sp.: KAMP, 1953: 61-62.
Distribution. U.S.A.: California (Lassen County).

5. *Grylloblatta chirurgica* GURNEY, 1961

GURNEY, 1961: 69, figs. 1-7. (Holotype: ♂, U.S.A., Washington, Skamania County, Ape Cave (1900 foot), in U.S. Nat. Mus., Washington, N 65377; allotype: ♀, some localities and dispositions).
KAMP, 1963: 65, fig. 43; 1970: 225, fig. 1; RENTZ, 1982: 5, 10.
Distribution. U.S.A.: S. Washington.

6. *Grylloblatta gurneyi* KAMP, 1963.

KAMP, 1963: 62, figs. 23-42. (Holotype: ♂, U.S.A., California, Modoc County, Merrill Ice Cave (4700 foot), in U.S. Nat. Mus., Washington; allotype: ♀, some localities and dispositions).
KAMP, 1970: 225, fig. 1; RENTZ, 1982: 5, 7, 10.
Distribution. U.S.A.: California (Modoc County).

7. *Grylloblatta occidentalis* SILVESTRI, 1931.

SILVESTRI, 1931: 293, figs. 1-3. (Pro var. *Grylloblatta campodeiformis*; holotype: ♂ (larvae), U.S.A., Washington, Mt. Baker (4200-5000 foot), presumably in Agricultural Zoology Laboratory, Portici, Italy or lost).
Grylloblatta campodeiformis var. *occidentalis*: BEAMER, 1933: 334; WALKER, 1937: 8-9.
Grylloblatta campodeiformis occidentalis: SILVESTRI, 1934: 18; GURNEY, 1937: 160, fig. 1 (brief description of imago ♂ and ♀ in key); 1948: 90 (part.), 98, figs. 4, 11, text-fig. B; 1953: 235, fig. 8a; KAMP, 1953: 61; 1963: 65, fig. 43; 1970: 225, fig. 1; RENTZ, 1982: 5, 10, 12.
Grylloblatta occidentalis: VICKERY and McE-KEVAN, 1983: 242, 250.
Distribution. U.S.A.: N. Washington (Mt. Baker and Hanagen Peak).

8. *Grylloblatta rothi* GURNEY, 1953.

GURNEY, 1953: 326, figs. 1, 5, 6, 10-12, 14, 16. (Holotype: ♂, U.S.A., Oregon, Happy Valley on Century Drive, about 15-20 miles S. Three Sisters (6450 foot), in U.S. Nat. Mus., Washington, N 61656; allotype: ♀, Oregon, Crater Lake).

KAMP, 1963: 65, fig. 43; RENTZ, 1982: 5, 7, 10.

Grylloblatta campodeiformis: ELSEA, 1937: 57.

Grylloblatta sp.: GURNEY, 1948: 90, fig. 11.

Grylloblatta rothi (lapsus calami): KAMP, 1970: 225, fig. 1.

Distribution. U.S.A.: S. Oregon.

9. *Grylloblatta scudderi* KAMP, 1979.

KAMP, 1979: 30, figs. 28-36. (Holotype: ♂, Canada, British Columbia, Garibaldi Provincial Park, Whistler Mountain (1951 m), in Canadian National Collection, Ottawa; allotype: ♀, some localities in Entomol. Mus. University of British Columbia).

RENTZ, 1982: 5, 10; VICKERY and McE-KEVAN, 1983: 242, 248.

Grylloblatta campodeiformis occidentalis: GURNEY, 1948: 90 (part.).

Distribution. Canada: S.W. British Columbia.

10. *Grylloblatta sculleni* GURNEY, 1937.

GURNEY, 1937: 161, 164, figs. 7-11. (Holotype: ♀, U.S.A., Oregon, Cascade Mountain, Three Sisters, Scott Camp (6600 foot), in U.S. Nat. Mus., Washington, N 52017).

GURNEY, 1948: 90, 98, fig. 11; 1953: 326, 330-331; KAMP, 1953: 61;

GURNEY, 1961: 72; KAMP, 1963: 57, 65, figs. 8, 9, 43; 1970: 225, fig. 1;

RENTZ, 1982: 5, 7, 10.

Distribution. U.S.A.: Oregon.

11. *Grylloblatta washoe* GURNEY, 1961.

GURNEY, 1961: 68, figs. 10-17. (Holotype: ♂, U.S.A., California, Eldorado County, Echo Summit on 4 miles S. Meyers, in American Mus. Nat. Hist., New York).

KAMP, 1963: 65, fig. 43; 1970: 225, fig. 1; RENTZ, 1982: 5, 10.

Distribution. U.S.A.: California (Eldorado County).

Populations of *Grylloblatta insertae sedis*.

There are 4 populations in U.S.A.: California, Idaho, Oregon and Washington (GURNEY, 1953: 330; 1961: 73; KAMP, 1963: 65; EDWARDS, 1982: 19) and 4 in Canada: British Columbia (VICKERY and McE-KEVAN, 1983: 250-251) known only by immature specimens and in all probability may be a new species.

2. Genus *Grylloblattina* BEY-BIENKO, 1951.

BEY-BIENKO, 1951: 507. Type species: *Grylloblattina djakonovi* BEY-BIENKO by original designation.

GURNEY, 1953: 331; ASAHINA, 1971: 202; RENTZ, 1982: 1.

1 species in USSR (Primorje).

1. *Grylloblattina djakonovi* BEY-BIENKO, 1951.

BEY-BIENKO, 1951: 507, figs. 1 (1-3), 2. (Holotype: ♀, USSR, Primorje, Island Petrov in Kievka Bay about 20 km W. Priobrashiniye, lost).

GURNEY, 1953: 331; ASAHINA, 1971: 202; PRAVDIN and STOROZHENKO, 1977: 353.

Distribution. USSR: Primorje (Island Petrov).

3. Genus *Galliosiana* CAUDELL, 1924.

CAUDELL, 1924a: 92 (nom. n. pro *Galliosia* CAUDELL et KING, preoccupied by HUSTACHE, 1920 in *Coleoptera*). Type species: *Galloisia nipponensis* CAUDELL et KING by original designation.

GURNEY, 1937: 160-161; 1948: 86-90, 98; FUKUSHIMA, 1969a: 16;

ASAHINA, 1971: 202; PRAVDIN and STOROZHENKO, 1977: 355;

RENTZ, 1982: 1; NAMKUNG, 1982: 32.

Galloisia CAUDELL et KING, 1924: 53.

Grylloblatta: SILVESTRI, 1927: 112 (part.).

2 subgenera, 10 species in Japan, S. Korea and USSR: Siberia (Altai) and Primorje.

3a. Subgenus *Galloisiana* CAUDELL, 1924.

Literature and type species see under genus.

1. *Galloisiana (Galloisiana) kiyosawai* ASAHINA, 1959.

ASAHINA, 1959: 250, figs. 3a, d. (Holotype: ♀, Central Japan, prefecture Gifu, Hirayu-onsen, in Tokyo Metropolitan University, Japan).

GURNEY, 1961: 75; FUKUSHIMA, 1969b: 16-18, figs. C (1-3) (♂): PRAVDIN and STOROZHENKO, 1977: 376.

Distribution. Central Japan: prefectures Nagano, Gifu and Chichibu.

2. *Galloisiana (Galloisiana) kosuensis* NAMKUNG, 1974.

Namkung, 1974b: 91, figs. 1-8 (Holotype: ♂, Korea, 480 km S.E. Seoul, Chung-cheong-buk-do, Danyong-gun, Kosudong-gul Cave at Kosu-ri in Daegang-myeon, in coll. NAMKUNG, Seoul; allotype: ♀, some localities and dispositions).

Namkung, 1982: 38-41, fig. 4.

Distribution. S. Korea: province Chung-cheong-buk-do.

3. *Galloisiana (Galloisiana) kurentzovi* PRAVDIN et STOROZHENKO, 1977 PRAVDIN et STOROZHENKO, 1977: 353, figs. 1-9. (Holotype: ♀, USSR, Primorje, National Park Kedrovaja Pad, in Zool. Institute USSR Acad. Sci., Leningrad; allotype: ♂, Mt. Krinitshnaja about 80 km N.E. Vladivostok, same disposition).

STOROZHENKO, 1979: 18-20; STOROZHENKO and OLEGER, 1984: 729.

Distribution. USSR: S. Primorje.

4. *Galloisiana (Galloisiana) nipponensis* (CAUDELL et KING, 1924).

CAUDELL et KING, 1924: 54, figs. 1-3. (*Galloisia*, holotype: ♂, Central Japan, Chuzenji near Nikko, in U.S. Nat. Mus., Washington, N 26848).

Galloisiana nipponensis: CAUDELL, 1924a: 92; SILVESTRI, 1927: 107-112; figs. I (1-13), II (1-8), III (1-4); KISHIDA, 1929: 104-108; MATSUMURA, 1931: 1373; ASAHINA and ASAHINA, 1933: 630; WALKER, 1937: 9; GURNEY, 1937: 160, figs. 2-6; 1948: 95-96, figs. 1, 2, 5-7 (♀); ASAHINA, 1959: 250, figs. 1a-d; GURNEY, 1961: 74-75; FUKUSHIMA, 1969b: 16-18, figs. A (1-3); PRAVDIN and STOROZHENKO, 1977: 356; NAGASHIMA, 1982: 52; YAMASAKI, 1982c: 13; 1982d: 333.

Galloisidea (sic.) *nipponensis*: SHIRAKI, 1932: 2038.

Distribution. Central Japan.

5. *Galloisiana (Galloisiana) pravdini* STOROZHENKO et OLIGER, 1984.

STOROZHENKO et OLIGER, 1984: 279, figs. 1-7 (Holotype: ♂, USSR, Altai, 100 km S.E. Gorno-Altai, in Zool. Institute USSR Acad. Sci., Leningrad; allotype: ♀, some locality and disposition).

Distribution. USSR: Altai.

6. *Galloisiana (Galloisiana) yasai* ASAHINA, 1959.

ASAHINA, 1959: 249, figs. 2a-d. (Holotype: ♂, Japan, prefecture Nagano, Tokudo-toge, in Tokyo Metropolitan University, Japan; allotype: ♀, prefecture Nagano, Shimajima-dani, some dispositions).

GURNEY, 1961: 75; FUKUSHIMA, 1969b: 16-18, figs. B (1-3).

Galloisiana yasai (sic!): PRAVDIN and STOROZHENKO, 1977: 355.

Distribution. Central Japan: prefectures Nagano, Gumma, Niigata and Chichibu.

7. *Galloisiana (Galloisiana) yezoensis* ASAHINA, 1961.

ASAHINA, 1961: 85, figs. 1-6. (Holotype: ♀, Japan, Hokkaido, Mt. Daisetsu, Miyozaki-toge near Yukomanbetsu, in Tokyo Metropolitan University, Japan).

FUKUSHIMA, 1969b: 16; PRAVDIN and STOROZHENKO, 1977: 355.

Distribution. Japan: Hokkaido.

3b. Subgenus *Ishiana* SILVESTRI, 1927.

SILVESTRI, 1927: 113 (pro subgenus *Grylloblatta*). Type species *Grylloblatta (Ishiana) notabilis* SILVESTRI, by original designation.

GURNEY, 1937: 160 (pro subgenus *Galloisiana*); 1948: 90; FUKUSHIMA, 1969a: 16; PRAVDIN and STOROZHENKO, 1977: 355.

8. *Galloisiana (Ishiana) biryongensis* NAMKUNG, 1974.

NAMKUNG, 1974a: 1-7, figs. 1-16. (Holotype: ♂, Korea, 500 km N.E. Seoul, Kangweon-do Province, Biryong-don-gul Cave at Biryong-dong of Yongtanri in Teongseon-myeon of Jengseon-gun, in coll. NAMKUNG; allotype: ♀, some localities and dispositions).

NAMKUNG, 1982: 34-38, figs. 2, 3.

Distribution. S. Korea: province Kangweon-do.

9. *Galloisiana (Ishiana) chuioi* GURNEY, 1961.

GURNEY, 1961: 73, figs. 18, 19; (Holotype: ♀, Japan, Kagawa prefecture, Island Megi-shima, Oninowaya Cave, in U.S. Nat. Mus., Washington, N 65378).

PRAVDIN and STOROZHENKO, 1977: 355.

Galloisiana sp.: Chujo, 1958: 1-7.

Distribution. Japan: prefecture Kagawa (island Megi-shima).

10. *Galloisiana (Ishiana) notabilis* (SILVESTRI, 1927).

SILVESTRI, 1927: 113-118, figs. IX, X (1-11), XI (1-4), XII (1-4). (*Grylloblatta (Ishiana)*, holotype: ♂ (larvae), Japan, Island Kyushu at Michion near Nagasaki, presumably in Agricultural Zoology Laboratory, Portici, Italy or lost).

Galloisiana (Ishiana) notabilis: GURNEY, 1937: 160; WALKER, 1937: 9; GURNEY, 1948: 96, 98; 1961: 74; PRAVDIN and STOROZHENKO, 1977: 355.

Galloisiana sp.: KAWASAWA, 1950: 1-2.

Distribution. Japan: island Kyushu.

Populations of *Galloisiana incertae sedis*.

There are about 25 populations in S.Korea (NAMKUNG, 1982: 30-32) and 3-4 in Japan (KAWASAWA, 1950: 1-2; FUKUSHIMA, 1969b: 18) known only by immature species mentioned and some of them may be new species.

Acknowledgments

The cooperation of the following persons, who have assisted me by making notes and some literature available, is gratefully acknowledged: Dr. H. ANDO, University of Tsukuba, Japan, Dr. S. ASAHINA, Tokyo, Japan,

Dr. G. FUKUSHIMA, Chichibu High School, Saitama, Japan, Dr. K. HARZ, Endsee, West Germany, Prof. D.K. McE. KEVAN, Lyman Entomol. Museum, McGill University, Canada, Dr. N. NAGASHIMA, Tokyo University of Agriculture, Japan, Prof. V.R. VICKERY, Lyman Entomol. Museum, Canada and Dr. T. YAMASAKI, National Sci. Museum, Tokyo, Japan.

Literature

- ANDO, H. and NAGASHIMA, T.: 1982. A preliminary note on the embryogenesis of *Galloisiana nipponensis* (CAUDELL et KING). In: Biology of the *Notoptera*, Japan: 89-95.
- ASAHINA, S. and E. ASAHINA: 1933. Insects fauna of Mt. Okuchichibu in Central Japan. *Konchyukai* I (6): 630-636 (in japan.).
- BACCETTI, B.: 1982. The Spermatozoon of *Arthropoda*. XXXII. *Galloisiana nipponensis* (CAUDELL et KING) (*Grylloblattodea*). In: Biology of the *Notoptera*, Japan: 71-78.
- BACCETTI, B.: 1981. Notulae Orthopterologicae. XXXII. Nuova notulae interpretazione dell'apparato riproduttore maschile nei *Grilloblattoidei*. *Mem. Soc. Entomol. Ital.* 60 (2): 49-60.
- BAKER, G.T.: 1982. Sensory receptors on the mandibles and labrum of *Grylloblatta campodeiformis* WALKER. *Zoologischer Anz.* 209 (5-6): 341-344.
- BEY-BIENKO, G.J.: 1962. On the general classification of Insects. *Rev. d'Entom. URSS.* 41 (1): 6-21 (in russian).
- BEY-BIENKO, G.J.: 1966. General Entomology. Moscow: 1-496 (in russian).
- BROWN, W.L.: 1982. *Grylloblattaria*. In: Parker S.P. (Ed.). Synopsis and classification of living organisms. Vol. 2. New York etc.: 351-352.
- BRUNER, L.: 1915. Preliminary catalogue of the *Orthopteroid* insects of the Philippine islands. *Univ. Studies Insects, Lincoln.* 15: 195-281.
- CAUDELL, A.N.: 1923b. Record of *Grylloblatta campodeiformis* in California. *Journ. Wash. Acad. Sci.* 13: 260-263.
- CORNER, J.A.: 1960: A further record of *Grylloblatta campodeiformis* WALKER in the interior of British Columbia. *Proc. Entomol. Soc. Brit. Columbia.* 57: 49.
- CRIDDLE, N.: 1926. The entomological record, 1925. 56th Ann. Rpt. Ent. soc. Ontario. 94-107.
- EDWARDS, J.G.: 1958. Exhibit of 2 color slides of *Grylloblatta barberi* at meeting of March 2, 1957. *Pan-Pacif. Ent.* 34 (1): 49.
- EYER, J.R.: 1924. The comparative morphology of the male genitalia of the primitive Lepidoptera. *Ann. Ent. Soc. Amer.*: 17 (3): 275-342.
- HEMING, B.S.: 1977. (Book review). MATSUDA, R. 1976. Morphology and evolution of the Insect Abdomen. *Questiones Ent.* 13 (2): 75-81.
- KAMP, J.M.: 1970. The cavernicolous *Grylloblattodea* of the western United States (1). *Ann. speleol.* 25 (1): 223-230.
- KEVAN, D.K. McE.: 1979. *Notoptera*. In: DANKS, H.V. (Ed.). Canada and its Insect Fauna. *Mem. Ent. Soc. Canada.* 108: 316-317.
- KENNEDY, C.H.: 1928. Evolutionary level in relation to geographic, seasonal and diurnal distribution of insects. *Ecology.* 9 (4): 367-379.
- LAMEERE, A.: 1935. *Precis de Zoologie.* Brussels.: 1-468.
- MAHOTIN, A.A.: 1952. Phylogeny of *Orthoptera* and morphology of its ovipositor. *Rev. d'Entomol. URSS.* 32: 126-136 (in russian).
- MAHOTIN, A.A.: 1953. Phylogeny of different groups of *Orthoptera* and morphology of ovipositor. *Mem. Inst. Animal Morphol., Moscow.* 8: 5-62 (in russian).
- MATSUDA, R.: 1965. Morphology and Evolution of the Insect Head. — *Mem. Amer. Entomol. Inst.* 4: 1-333.
- MATSUDA, R.: 1970. Morphology and evolution of the Insect Thorax. *Mem. Ent. Soc. Canada.* 76: 1-431.
- MATSUDA, R.: 1976. Morphology and evolution of the Insect Abdomen. Oxford, Pergamon press: 1-534.
- MATSUMURA, S.: 1931. 6000 Illustrated Insects of Japanese Empire. Tokyo: 1-1719 (in japan).
- MATSUZAKI, M., ANDO, H. and S.N. VISSCHER: 1982. Fine structure of Oocyte and Follicular Cells during Oogenesis in *Galloisiana nipponensis* (CAUDELL et KING). In: Biology of the *Notoptera*, Japan: 79-87.
- McIVER, S.B. and SUTCLIFFE, J.F.: 1982. Sensilla of *Grylloblatta campodeiformis* WALKER. *Ibid.*: 137-157.
- NAGASHIMA, T.: 1982a. Anatomy of *Galloisiana nipponensis* (CAUDELL et KING). *Ibid.*: 113-135.
- NAGASHIMA, T., ANDO, H. and G. FUKUSHIMA: 1982. Life history of *Galloisiana nipponensis* (CAUDELL et KING). *Ibid.*: 43-59.
- NAMKUNG, J.: 1982. The taxonomy of the *Grylloblattodea* and their Distribution in Korea. *Ibid.*: 29-42.
- RÄHLE, W.: 1970. Untersuchungen an Kopf und Prothorax von *Embia ramburi* RIMSKY-KORSAKOW 1906 (*Embioptera, Embiidae*). *Zool. Jb. Anat.* 87: 248-330.
- RENTZ, D.C.F.: 1982. A review of the Systematics, Distribution and biometrics of the North American *Grylloblattidae*. In: Biology of the *Notoptera*, Japan.: 1-18.
- ROHDENDORF, B.B.: 1977. On the codification of names of higher taxa in zoology. *Paleontol. Journ.* 2: 14-22 (in russian).
- SHIRAKI, T.: 1932. *Orthoptera*. In: *Iconographia Insectorum Japonicorum.* Tokyo: 2024-2114 (in Japanese).
- SILVESTRI, F.: 1934. *Compendio di Entomologica Applicata. Parte Speciale*, vol. 1, Parte 1: 1-448.
- SPENCER, G.J.: 1952. The status of our knowledge of the insects of British Columbia. *Proc. Ent. Soc. Brit. Columbia,* 48: 36-41.

- STOROZHENKO, S.Yu.: 1979. Habitat and behavior of *Galloisiana kurentzovi* PRAVDIN et STOROZHENKO (*Insecta, Grylloblattida*) in southern Primorje Territory. Biol. Sci., Moscow. 2: 18-21 (in russian).
- STOROZHENKO, S.Yu and A.I. OLIGER: 1984. A new species of *Grylloblattida* from North-East Altai. Rev. d'Entom. URSS. 63 (4): 729-732 (in russian).
- VICKERY, V.R. and JOHNSTON, D.E. and D.K. McE. KEVAN: 1974. The orthopteroid insects of Quebec and the Atlantic provinces of Canada. Mem. Lyman. Ent. Mus. Res. Lab. 1: 1-204.
- VICKERY, V.R. and D.K. McE KEVAN: 1983. A monograph of the *Orthopteroid* insects of Canada and adjacent regions. Vol. 1. Quebec: 1-679.
- WALKER, E.M.: 1920. The life cycle of the orthopteroid orders. Ann. Ent. Soc. Amer. 13: 137-141.
- YAMASAKI, T.: 1982a. Bibliography on the *Notoptera*. In: Biology of the *Notoptera*, Japan: 173-181.
- YAMASAKI, T.: 1982b. Nymphal morphology in early postembryonic stages of *Galloisiana nipponensis* (CAUDELL et KING). Ibid.: 97-111.
- YAMASAKI, T.: 1982d. The *Orthoptera* and *Notoptera* of Ozegahara Moor. In: Ozegahara: Scientific Researches of the Highmoor in Central Japan. Tokyo: 331-334.

Address of author:

Dr. S. Yu. STOROZHENKO,
Institut of Biology and Pedology
Far East Science Centre, USSR Academy of Sciences
Vladivostok, 690022, USSR

This excellent catalogus on the present *Grylloblattida* will in its arrangement be a great help to all who are working with them or intend to study them. Exact references to the books and journals in which the mentioned publications are published you find in T. YAMASAKI: 1982 in „Biology of the *Notoptera*“ (edited by HIROSHI ANDO) Kashigo-insatsu Co. Ltd., Nagano, Japan.

Kurt HARZ